

Sources:

1. [Royal College of Physicians Sentinel Stroke National Audit Programme \(2016\) National clinical audit annual results portfolio March 2016-April 2017](#)
2. Saka, O, McGuire A, Wolfe, C. Cost of stroke in the United Kingdom. *Age and Ageing* 2009;38:27–32.
3. Healey J.S, Connolly S.J, Gold, M.R. Subclinical atrial fibrillation and the risk of stroke. *New England Journal of Medicine* 2012;366:120-129.
4. Borowsky L.H, Regan S, Chang Y et al. First diagnosis of atrial fibrillation at the time of stroke. *Cerebrovascular Disease*. 2017;43(3–4):192–199.
5. Björck S, Atrial fibrillation, stroke risk, and warfarin therapy revisited: a population-based study. *Stroke* 2013; 44:3103–3108.
6. [Public Health England: Atrial fibrillation prevalence estimates in England: application of recent population estimates of AF in Sweden. England: Public Health England 2015.](#)
7. Odutayo A, Wong C.X, Hsiao A.J et al. Atrial fibrillation and risks of cardiovascular disease, renal disease, and death: systematic review and meta-analysis. *BMJ* 2016;354:i4482.
8. Krijthe B.P, Kunst A, Benjamin E.J et al. Projections on the number of individuals with atrial fibrillation in the European Union, from 2000 to 2060. *European Heart Journal* 2013;34: 2746–2751.
9. [Nice Quality standards for AF](#)